Canadian Journal of Anesthesia

Index of Authors

Volume 49, December 2002

Note: Abstracts and refresher course outlines in the May *Annual Meeting Supplement* are not included in this index.

A

Abdallah C, Karsli C, Bissonnette B. Fentanyl is more effective than remifentanil at preventing increases in cerebral blood flow velocity during intubation in children, 1070

Aboulcish E. See *Ezri T,* jt. auth.

Agarwal A, Bose N, Gaur A, Singh U, Gupta MK, Singh D. Acupressure and ondansetron for postoperative nausea and vomiting after laparoscopic cholecystectomy, 554

Agarwal A, Singh PK, Dhiraj S, Pandey CM, Singh U. Oxygen in air ($FI0_2$ 0.4) improves gas exchange in young healthy patients during general anesthesia, 1040

Agrò F, Antonelli S, Cataldo R, Montecchia F, Barzoi G, Petitti T. The ProSeal laryngeal mask airway: fibreoptic visualization of the glottic opening is associated with ease of insertion of the gastric tube, 867

Ahmed MZ, Vohra A. The reinforced laryngeal mask airway (RLMA) protects the airway in patients undergoing nasal surgery - an observational study of 200 patients, 863

Ainsworth L. See *Pollard RC,* jt. auth.

Al-Metwalli R, Mowafi HA. A modification of the inter-cuff technique of IVRA for use in knee arthroscopy, 687

Aldwinckle RJ, Carr AS. The anesthetic management of a patient with Emery-Dreifuss muscular dystrophy for orthopedic surgery, 467

Alkaissi A, Evertsson K, Johnsson V-A, Ofenbartl L, Kalman S. P6 acupressure may relieve nausea and vomiting after gynecological surgery: an effectiveness study in 410 women, 1034

Alsuwaida K. Primary cutaneous mucormycosis complicating the use of adhesive tape to secure the endotracheal tube, 880

Antonelli M. See *Cavaliere F,* jt. auth.*; Conti G,* jt. auth.

Antonelli S. See *Agrò F,* jt. auth.

Antoniou T. See *Theodoraki K,* jt. auth.

Aoyama K. See *Takenaka I,* jt. auth.

Aoyama Y. See *Arakawa M,* jt. auth.

Arakawa M, Aoyama Y, Ohe Y. Epidural bolus injection with alkalinized lidocaine improves blockade of the first sacral segment - a brief report, 566

Arcangeli A. See *Cavaliere F,* jt. auth.

Archer DP, McTaggart Cowan RA, Falkenstein RJ, Sutherland GR. Intraoperative mobile magnetic resonance imaging for craniotomy lengthens the procedure but does not increase morbidity, 420

Argalious M. See *Farag E,* jt. auth.

Armstrong KP, Kennedy B, Watson JT, Morley-Forster PK, Yee I, Butler R. Epinephrine reduces the sedative side effects of epidural sufentanil for labour analgesia, 72

Arndt GA. See *Yun ES,* jt. auth.

Arriero MM, Muñoz Alameda L, López-Farré A, Escribano Burgos M, Carrasco C, Millás I, Celdrán A, Pinta JC de la. Sevoflurane reduces endothelium-dependent vasorelaxation: role of superoxide anion and endothelin, 471

Atanassoff PG, Hartmannsgruber MWB. Central nervous system side effects are less important after iv regional anesthesia with ropivacaine 0.2% compared to lidocaine 0.5% in volunteers, 169

Atef A. See *Elhakim M,* jt. auth.

Aya G. See *Lefrant J-Y,* jt. auth.

Aye T, Milne B. Ketamine anesthesia for pericardial window in a patient with pericardial tamponade and severe COPD, 283

B

Babin D. See *Couture P,* jt. auth.*; Denault AY,* jt. auth.*; Hemmerling TM,* jt. auth.*; Taillefer J,* jt. auth.

Badaoui R, El Kettani C, Ouendo M, Fikri M, Montravers P. Bilateral pneumothorax following tracheal extubation (Images in anesthesia), 90

Baenziger O. See *Weiss M,* jt. auth.

Bahk J-H. See *Park H-P,* jt. auth.

Baker A. See *White H,* jt. auth.

Balladur P. See *Beaussier M,* jt. auth.

Balmer C. See *Weiss M,* jt. auth.

Bands CJ. See *Laupland KB,* jt. auth.

Baraka A, Siddik-Sayyid S, Jalbout M, Yaacoub C. Variable hemodynamic fluctuations during resection of multicentric extraadrenal pheochromocytomas, 682

Barat M. See *Sesay M*, jt. auth.
Barzoi G. See *Agrò F*, jt. auth.
Batra YK. See *Mahajan R*, jt. auth.
Baxter FJ, Randall J, Miller JD, Higgins DA, Powles ACP, Choi PT-L. Rescue therapy with inhaled nitric oxide in critically ill patients with severe hypoxemic respiratory failure (brief report), 315
Beattie WS. See *Bradley J*, jt. auth.
Beaulieu M-A. See *Wherrett CG*, jt. auth.
Beaulieu P. See *Hemmerling TM*, jt. auth.
Beaulieu Y. See *Denault A*, jt. auth.
Beaussier M, Decorps A, Tilleul P, Megnigbeto A, Balladur P, Lienhart A. Desflurane improves the throughput of patients in the PACU. A cost-effectiveness comparison with isoflurane, 339
Beh T, Splinter W, Kim J. In children, nitrous oxide decreases pain on injection of propofol mixed with lidocaine, 1061
Bélisle S. See *Denault A*, jt. auth.
Bell EA, Jones BP, Olufolabi AJ, Dexter F, Phillips-Bute B, Greengrass RA, Penning DH, Reynolds JD, Duke Women's Anesthesia Research Group. Iliohypogastric-ilioinguinal peripheral nerve block for post-Cesarean delivery analgesia decreases morphine use but not opioid-related side effects, 694
Benais H. See *Pierre S*, jt. auth.
Benhamou D, Técsy M, Parry N, Mercier FJ, Burg C. Audit of an early feeding program after Cesarean delivery: patient wellbeing is increased, 814
Bennett HL. See *Moskowitz DM*, jt. auth.
Benzer A. See *Lorenz IH*, jt. auth.
Beriault MT. See *Maltby JR*, jt. auth.
Bettex DA, Schmidlin D, Chassot P-G, Schmid ER. Intrathecal sufentanil-morphine shortens the duration of intubation and improves analgesia in fast-track cardiac surgery, 711
Bhardwaj V. See *Singh B*, jt. auth.
Bissonnette B, Paut O. Active warming of saline or blood is ineffective when standard infusion tubing is used: an experimental study, 270
- see also *Abdallah C*, jt. auth.; *Karsli C*, jt. auth.
Blanshard H. See *Wong J*, jt. auth.
Boker A, Brownell L, Donen N. The Amsterdam preoperative anxiety and information scale provides a simple and reliable measure of preoperative anxiety, 792
Boldt J. See *Piper SN*, jt. auth.
Borgeat A. See *Ekatodramis G*, jt. auth.
Bose N. See *Agarwal A*, jt. auth.; *Pandey CK*, jt. auth.
Boudreault D. See *Denault AY*, jt. auth.
Boulard G. See *Sesay M*, jt. auth.
Boyer D. See *Smith CE*, jt. auth.
Bracco D, Parlow JL. Best evidence in anesthetic practice (prevention): Dopamine does not prevent death, acute renal failure, or need for dialysis; commentaries, 417
Bradley J, McAlister FA, McRae K, Beattie WS. Best evidence in anesthetic practice: Clinical prediction guide: a 14-item index predicts 30-day risk of postoperative pneumonia after non-cardiac surgery; commentaries, 655
Brimacombe J, Keller C, Roth W, Loeckinger A. Large cuff volumes impede posterior pharyngeal mucosal perfusion with the laryngeal tube airway, 1084
Brownell L. See *Boker A*, jt. auth.
Bruelle P. See *Lefrant J-Y*, jt. auth.
Bruskin A. See *Kligman M*, jt. auth.
Bryson G, Choi P. Best evidence in anesthetic practice: Albumin neither increases nor decreases mortality in critically ill patients; commentaries, 620
Bucher H-U. See *Weiss M*, jt. auth.
Buckenmaier CC III, Steele SM, Nielsen KC, Klein SM. Paravertebral somatic nerve blocks for breast surgery in a patient with hypertrophic obstructive cardiomyopathy, 571
Buithieu J. See *Denault AY*, jt. auth.
Burg C. See *Benhamou D*, jt. auth.
Burjorjee JE, Milne B. Propofol for electrical storm; a case report of cardioversion and suppression of ventricular tachycardia by propofol, 973
Butler R, Laufer B. Best evidence in anesthetic practice (prevention): recombinant human activated protein C reduces mortality in severe sepsis; commentaries, 207
- see also *Armstrong KP*, jt. auth.
Byrick R. See *Doyle DJ*, jt. auth.; *Jacka MJ*, jt. auth.
Byrick RJ, Craig D, Carli F. A physician workforce planning model applied to Canadian anesthesiology: assessment of needs, 663
- see also *Craig D*, jt. auth.

C

Caillé J-M. See *Sesay M*, jt. auth.
Campbell N, Rosaeg OP. Anesthetic management of a parturient with Ehlers Danlos syndrome type IV, 493
- see also *Rosaeg OP*, jt. auth.
Capt H. See *Pradervand Mooser M*, jt. auth.
Carli F. See *Byrick RJ*, jt. auth.; *Craig D*, jt. auth.
Carr AS. See *Aldwinckle RJ*, jt. auth.
Carrasco C. See *Arriero MM*, jt. auth.
Cartier R. See *Couture P*, jt. auth.
Cartotto RC. See *Cooper AB*, jt. auth.
Cashin F. See *Doyle DJ*, jt. auth.
Cataldo R. See *Agrò F*, jt. auth.
Cathrein P. See *Ekatodramis G*, jt. auth.

Caton D. See *Frölich MA*, jt. auth.
Cavaliere F, Antonelli M, Arcangeli A, Conti G, Costa R, Pennisi MA, Proietti R. A low-dose remifentanil infusion is well tolerated for sedation in mechanically ventilated, critically-ill patients, 1088
Celdrán A. See *Arriero MM*, jt. auth.
Chan VWS. See *Salomons TV*, jt. auth.
Chand MB. See *Shrestha BM*, jt. auth.
Chang D-P. See *Pang W-W*, jt. auth.
Chang H, Tsai S-Y, Chang Y, Chen T-L, Chen R-M. Therapeutic concentrations of propofol protects mouse macrophages from nitric oxide-induced cell death and apoptosis, 477
Chang Y. See *Chang H*, jt. auth.
Chang Y-C. See *Wong C-S*, jt. auth.
Chassot P-G. See *Bettex DA*, jt. auth.
Chen L-K. See *Huang Y-C*, jt. auth.
Chen R-M. See *Chang H*, jt. auth.
Chen RJB. See *Joo HS*, jt. auth.
Chen T-L. See *Chang H*, jt. auth.
Chen X. See *Yamakage M*, jt. auth.
Cherng C-H. See *Lu C-H*, jt. auth.; *Wong C-S*, jt. auth.
Chia Y-Y, Chow L-H, Hung C-C, Liu K, Ger L-P, Wang P-N. Gender and pain upon movement are associated with the requirements for postoperative patient-controlled *iv* analgesia: a prospective survey of 2,298 Chinese patients, 249
Choi P. See *Bryson G*, jt. auth.
Choi PT-L, Galinski SE, Lucas S, Takeuchi L, Jadad AR. Examining the evidence in anesthesia literature: a survey and evaluation of obstetrical postdural puncture headache reports, 49
- see also *Baxter FJ*, jt. auth.; *Smith KJ*, jt. auth.
Chow L-H. See *Chia Y-Y*, jt. auth.
Chowvanayotin S. See *Kongsayreepong S*, jt. auth.
Choyce A, Peng P. A systemic review of adjuncts for intravenous regional anesthesia for surgical procedures (review article), 32
Chung F. See *Imasogie N*, jt. auth.; *Jenkins K*, jt. auth.; *Rocchi A*, jt. auth.; *Wong J*, jt. auth.
Chung PCH, Li AH, Lin CC, Yang MW. Elevated vascular resistance after labetalol during resection of a pheochromocytoma (brief report), 148
Cina C. See *Fayad A*, jt. auth.
Cleave-Hogg D. See *Morgan PJ*, jt. auth.
Cohen MM. See *Jacka MJ*, jt. auth.
Cohendy R. See *Lefrant J-Y*, jt. auth.
Colagiovanni D. See *Favaro R*, jt. auth.
Comunale ME. See *Park KW*, jt. auth.
Connelly NR. See *Parker RK*, jt. auth.
Conti G, De Cosmo G, Grazia Bocci M, Antonelli M, Ferro G, Costa R, Zito G, Proietti R. Alfentanil does not increase resistance of the respiratory system in ASA I patients ventilated mechanically during general anesthesia, 718
- see also *Cavaliere F*, jt. auth.
Cook D. See *Skrobik Y*, jt. auth.
Cooper AB, Islur A, Gomez M, Goldenson GL, Cartotto RC. Hypercapnic respiratory failure and partial upper airway obstruction during high frequency oscillatory ventilation in an adult burn patient, 724
Correa R. See *Jenkins K*, jt. auth.
Costa R. See *Cavaliere F*, jt. auth.; *Conti G*, jt. auth.
Cousineau KM. See *Moskowitz DM*, jt. auth.
Couture P, Denault A, Limoges P, Sheridan P, Babin D, Cartier R. Mechanisms of hemodynamic changes during off-pump coronary artery bypass surgery, 835
- see also *Denault AY*, jt. auth.; *Taillefer J*, jt. auth.
Crago R. See *Imasogie N*, jt. auth.
Craig D, Byrick R, Carli F. A physician workforce planning model applied to Canadian anesthesiology: planning the future supply of anesthesiologists, 671
- see also *Byrick R*, jt. auth.
Critchley LAH. See *Hui JKL*, jt. auth.
Crosby E. Airway management after upper cervical spine injury: what have we learned?, 733
Crossan ML. See *Rosaeg OP*, jt. auth.
Crozat P. See *Sesay M*, jt. auth.
Cuenca-Peña J. See *Prieto-Alvarez P*, jt. auth.
Cunningham AJ. See *Duggan M*, jt. auth.
Czeslick E. See *Sablotzki A*, jt. auth.

D

Dahaba AA, Oettl K, von Klobucar F, Reibnegger G, List WF. End-stage renal failure reduces central clearance and prolongs the elimination half life of remifentanil, 369
Dai HB. See *Park KW*, jt. auth.
Dash NR. See *Pandey CK*, jt. auth.
Dauzat M. See *Lefrant J-Y*, jt. auth.
Davey R. See *Jenkins K*, jt. auth.
Davis JA. See *Jahr JS*, jt. auth.
De Cosmo G. See *Conti G*, jt. auth.
de Rossi L, Gott K, Horn N, Hecker K, Hutschenreuter G, Rossaint R. Xenon preserves neutrophil and monocyte function in human whole blood, 942
de Vicente J. See *Reinoso-Barbero F*, jt. auth.
DeBoer GE. See *Farag E*, jt. auth.
Decorps A. See *Beaussier M*, jt. auth.
Dehne MG. See *Sablotzki A*, jt. auth.
Demaria R. See *Lefrant J-Y*, jt. auth.
Denault A, Beaulieu Y, Bélisle S, Peachey G. Best evi-

dence in anesthetic practice (treatment): vasopressin neither improves nor worsens survival from cardiac arrest; commentaries, 312
- see also *Couture P*, jt. auth.; *Taillefer J*, jt. auth.
Denault AY, Couture P, McKenty S, Boudreault D, Plante F, Perron R, Babin D, Buithieu J. Perioperative use of transesophageal echocardiography by anesthesiologists: impact in noncardiac surgery and in the intensive care unit, 287
Deng X-M. See *Xiao W-J*, jt. auth.
Devitt JH. See *Jacka MJ*, jt. auth.
Dexter F. See *Bell EA*, jt. auth.
Dhiraj S. See *Agarwal A*, jt. auth.
Di Lascio F. See *Favaro R*, jt. auth.
Dinesen TRJ. See *Ling E*, jt. auth.
Dixon K. See *Parker RK*, jt. auth.
Dobranowski J. See *Smith KJ*, jt. auth.
Doi K. See *Yamada Y*, jt. auth.
Donati F. See *Dragne A*, jt. auth.; *Hemmerling TM*, jt. auth.; *Motamed C*, jt. auth.
Donen N. See *Boker A*, jt. auth.
Douglas MJ, Swenerton JE. Epidural anesthesia in three parturients with lumbar tattoos: a review of possible implications, 1057
Dousset V. See *Sesay M*, jt. auth.
Dowd N. See *Duggan M*, jt. auth.
Doyle DJ, Byrick R, Filipovic D, Cashin F. Silica zeolite scavenging of exhaled isoflurane: a preliminary report, 799
Doyle J. See *Joo HS*, jt. auth.
Dragne A, Varin F, Plaud B, Donati F. Rocuronium pharmacokinetic-pharmacodynamic relationship under stable propofol and isoflurane anesthesia, 353
Driessen B. See *Jahr JS*, jt. auth.
Duggan M, Dowd N, O'Mara D, Harmon D, Tormey W, Cunningham AJ. Benzodiazepine premedication may attenuate the stress response in daycase anesthesia: a pilot study, 932
- see also *O'Brien B*, jt. auth.
Duke PC. See *Lambert AS*, jt. auth.
Duke Women's Anesthesia Research Group. See *Bell EA*, jt. auth.
Dullenkopf A, Holzmann D, Feurer R, Gerber A, Weiss M. Tracheal intubation in children with Morquio syndrome using the angulated videointubation laryngoscope, 198
Dunn SM. See *Parker RK*, jt. auth.
Duvaldestin P. See *Slavov V*, jt. auth.
DuVall D. See *Ling E*, jt. auth.

E

Ecoffey C. See *Estèbe J-P*, jt. auth.; *Langlois G*, jt. auth.
Egawa H, Minami J, Fujii K, Hamaguchi S, Okuda Y, Kitajima T. QT interval and QT dispersion increase in the elderly during laparoscopic cholecystectomy: a preliminary study, 805
Eidelman LA. See *Orbach-Zinger S*, jt. auth.
Ekatodramis G, Min K, Cathrein P, Borgeat A. Use of a double epidural catheter provides effective postoperative analgesia after spine deformity surgery, 173
El Kettani C. See *Badaoui R*, jt. auth.
El-Mansouri M. See *Parker RK*, jt. auth.
Eledjam J-J. See *Lefrant J-Y*, jt. auth.
Elhakim M, Nafie M, Mahmoud K, Atef A. Dexamethasone 8 mg in combination with ondansetron 4 mg appears to be the optimal dose for the prevention of nausea and vomiting after laparoscopic cholecystectomy, 922
Erez I. See *Shenkman Z*, jt. auth.
Ergin MA. See *Moskowitz DM*, jt. auth.
Escribano Burgos M. See *Arriero MM*, jt. auth.
Esposito G. See *Favaro R*, jt. auth.
Estèbe J-P, Gentili ME, Le Corre P, Le Verge R, Moulinoux J-P, Ecoffey C. Sciatic nerve block with bupivacaine-loaded microspheres prevents hyperalgesia in an inflammatory animal model, 690
- see also *Langlois G*, jt. auth.
Etchells EE. See *Jenkins K*, jt. auth.
Evans NR, Llewellyn R, Gardner SV, James MFM. Aspiration prevented by the ProSeal™ laryngeal mask airway: a case report, 413
Evertsson K. See *Alkaissi A*, jt. auth.
Evron S. See *Ezri T*, jt. auth.
Ezri T, Abouleish E, Lee C, Evron S. Intracranial subdural hematoma following dural puncture in a parturient with HELLP syndrome, 820

F

Faheem U. See *Parker RK*, jt. auth.
Falkenstein RJ. See *Archer DP*, jt. auth.
Fanning A. See *Salomons TV*, jt. auth.
Farag E, Argalious M, Narouze S, DeBoer GE, Tome J. The anesthetic management of ventricular septal defect (VSD) repair in a child with mitochondrial cytopathy, 958
Farley FA. See *Munro HM*, jt. auth.
Farrar M. See *Karsli C*, jt. auth.
Fasnacht M. See *Weiss M*, jt. auth.
Fasting S, Gisvold SE. Serious intraoperative problems - a five-year review of 83,844 anesthetics, 545
Favaro R, Tordiglione P, Di Lascio F, Colagiovanni D, Esposito G, Quaranta S, Gasparetto A. Effective nasotracheal intubation using a modified transillumination technique, 91

Fayad A, Sawchuk C, Yang H, Cina C. Transesophageal echocardiography in the management of left atrio-femoral bypass during thoracoabdominal aortic aneurysm repair: a case report, 1081
Fellmann C. See *Neff TA*, jt. auth.
Ferro G. See *Conti G*, jt. auth.
Feurer R. See *Dullenkopf A*, jt. auth.
Fick GH. See *Maltby JR*, jt. auth.
Fikri M. See *Badaoui R*, jt. auth.
Filep JG. See *Skrobik YK*, jt. auth.
Filipovic D. See *Doyle DJ*, jt. auth.
Fischer JE. See *Weiss M*, jt. auth.
Flynn N. See *O'Brien B*, jt. auth.
Foëx P. See *Kato R*, jt. auth.
Forte L. See *Rocchi A*, jt. auth.
Freud E. See *Shenkman Z*, jt. auth.
Friedrich I. See *Sablotzki A*, jt. auth.
Frölich MA, Caton D. Baseline heart rate may predict hypotension after spinal anesthesia in prehydrated obstetrical patients, 185
Fuechslin RM. See *Neff TA*, jt. auth.
Fujii K. See *Egawa H*, jt. auth.
Fujii Y, Toyooka H. High-dose colforsin daropate increases diaphragmatic contractility in dogs, 877
Fukuyama H. See *Takenaka I*, jt. auth.

G

Galinski SE. See *Choi PT-L*, jt. auth.
Gardaz J-P. See *Pradervand Mooser M*, jt. auth.
Gardner SV. See *Evans NR*, jt. auth.
Gasparetto A. See *Favaro R*, jt. auth.
Gaur A. See *Agarwal A*, jt. auth.
Gautam MP. See *Shrestha BM*, jt. auth.
Gentili ME. See *Estève J-P*, jt. auth.; *Langlois G*, jt. auth.
Ger L-P. See *Chia Y-Y*, jt. auth.
Gerber A. See *Dullenkopf A*, jt. auth.
Gerber AC. See *Weiss M*, jt. auth.
Gerberc AC. See *Neff TA*, jt. auth.
Ghosh BR. See *Manjushree R*, jt. auth.
Gibson C. See *Parker RK*, jt. auth.
Gin T. See *Lee A*, jt. auth.
Girard A. See *Taillefer J*, jt. auth.
Gisvold SE. See *Fasting S*, jt. auth.
Godec M. See *Vintar N*, jt. auth.
Goldenson GL. See *Cooper AB*, jt. auth.
Goldstein DH. See *VanDenKerkhof EG*, jt. auth.
Gomar-Sancho C. See *Prieto-Alvarez P*, jt. auth.
Gómez-Criado MS. See *Reinoso-Barbero F*, jt. auth.
Gomez M. See *Cooper AB*, jt. auth.
Goobiec SM. See *Pollard RC*, jt. auth.
Gosselin R. See *Jahr JS*, jt. auth.
Gott K. See *de Rossi L*, jt. auth.
Grady D. See *Wong J*, jt. auth.
Grazia Bocci M. See *Conti G*, jt. auth.
Greengrass RA. See *Bell EA*, jt. auth.
Gritznig I. See *Lorenz IH*, jt. auth.
Grond S. See *Sablotzki A*, jt. auth.
Guerin J. See *Sesay M*, jt. auth.
Gunnaleka P. See *Kongsayreepong S*, jt. auth.
Gunther RA. See *Jahr JS*, jt. auth.
Gupta MK. See *Agarwal A*, jt. auth.
Gutermacher M. See *Shenkman Z*, jt. auth.

H

Hall R. See *Murkin J*, jt. auth.
Halpern S. See *Macarthur A*, jt. auth.
Ham B-M. See *Park H-P*, jt. auth.
Hamaguchi S, Okuda Y, Kitajima T, Masawa N. Five percent, 7.5% or 10% hypertonic saline prevents delayed neuronal death in gerbils, 745
- see also *Egawa H*, jt. auth.
Han S-M. See *Lee E-J*, jt. auth.
Handa K. See *Manjushree R*, jt. auth.
Harding G. See *Hudson RJ*, jt. auth.
Harmon D. See *Duggan M*, jt. auth.; *O'Brien B*, jt. auth.
Hartmannsgruber MWB. See *Atanassoff PG*, jt. auth.
Hecker K. See *de Rossi L*, jt. auth.
Hemmerling TM, Donati F, Babin D, Beaulieu P. Duration of control stimulation does not affect onset and offset of neuromuscular blockade at the corrugator supercilii muscle measured with phonomyography or acceleromyography, 913
- see also *Jonas JB*, jt. auth.
Henderson BT. See *Hudson RJ*, jt. auth.
Hentschel T. See *Sablotzki A*, jt. auth.
Hervilla S. See *Reinoso-Barbero F*, jt. auth.
Higgins DA. See *Baxter FJ*, jt. auth.
Ho AM-H, Karmakar MK. Combined paravertebral lumbar plexus and parasacral sciatic nerve block for reduction of hip fracture in a patient with severe aortic stenosis, 946
Ho S-T. See *Lu C-H*, jt. auth.
Holzmann D. See *Dullenkopf A*, jt. auth.
Hoppenstein D. See *Shenkman Z*, jt. auth.
Horn N. See *de Rossi L*, jt. auth.
Huang C-H. See *Huang Y-C*, jt. auth.
Huang G-S. See *Wong C-S*, jt. auth.
Huang M-H. See *Pang W-W*, jt. auth.
Huang Y-C, Tsai S-K, Huang C-H, Wang M-H, Lin P-L, Chen L-K, Lin C-J, Sun W-Z. Intravenous tenoxicam reduces uterine cramps after Cesarean delivery, 384
Hudson RJ, Thomson IR, Henderson BT, Singh K, Harding G, Peterson DJ. Validation of fentanyl pharmacokinetics in patients undergoing coronary artery bypass grafting, 388

Hui JKL, Critchley LAH, Karmakar MK, Lam PKK. Co-administration of alfentanil-propofol improves laryngeal mask airway insertion compared to fentanyl-propofol, 508
Hung C-C. See *Chia Y-Y*, jt. auth.
Hutschenreuter G. See *de Rossi L*, jt. auth.
Hwang J-H. See *Lee E-J*, jt. auth.

I
Ichinose H. See *Nakayama M*, jt. auth.
Imasogie N, Crago R, Leyland NA, Chung F. Probable gas embolism during operative hysteroscopy caused by products of combustion, 1044
Inomata S. See *Miyabe M*, jt. auth.
Islur A. See *Cooper AB*, jt. auth.
Itano Y. See *Yokoyama M*, jt. auth.

J
Jacka MJ, Cohen MM, To T, Devitt JH, Byrick R. The appropriateness of the pulmonary artery catheter in cardiovascular surgery, 276
Jadad AR. See *Choi PT-L*, jt. auth.
Jäger M. See *Jonas JB*, jt. auth.
Jahr JS, Lurie F, Driessen B, Davis JA, Gosselin R, Gunther RA. The HemoCue®, a point of care B-hemoglobin photometer, measures hemoglobin concentrations accurately when mixed *in vitro* with canine plasma and three hemoglobin-based oxygen carriers (HBOC), 243
Jain K. See *Mahajan R*, jt. auth.
Jalbout M. See *Baraka A*, jt. auth.
James MFM. See *Evans NR*, jt. auth.
Jedeikin R. See *Shenkman Z*, jt. auth.
Jenkins K, Chung F, Wennberg R, Etchells EE, Davey R. Fat embolism syndrome and elective knee arthroplasty, 19
Jenkins K, Wong DT, Correa R. Management choices for the difficult airway by anesthesiologists in Canada, 850
Johnson D, Truman C. Hospital practice more than specialty influences the choice of regional or general anesthesia for Cesarean section, 954
Johnsson V-A. See *Alkaissi A*, jt. auth.
Jonas JB, Jäger M, Hemmerling TM. Continuous retrobulbar anesthesia for scleral buckling surgery using an ultra-fine spinal anesthesia catheter, 487
Jones BP. See *Bell EA*, jt. auth.
Joo HS, Kataoka MT, Chen RJB, Doyle J, Mazer CD. PVC tracheal tubes exert forces and pressures seven to ten times higher than silicone or armoured tracheal tubes - an *in vitro* study, 986
Juan J-M. See *Lefrant J-Y*, jt. auth.

K
Kadoya T. See *Takenaka I*, jt. auth.
Kakiuchi Y. See *Miyabe M*, jt. auth.
Kalman S. See *Alkaissi A*, jt. auth.
Kamada Y. See *Yamakage M*, jt. auth.
Kamasumadram R. See *Parker RK*, jt. auth.
Kanaya N. See *Nakayama M*, jt. auth.
Karasawa F. See *Takamatsu I*, jt. auth.
Karmakar MK. See *Ho AM-H*, jt. auth.; *Hui JKL*, jt. auth.
Karsli C, Luginbuehl I, Farrar M, Bissonnette B. Propofol decreases cerebral blood flow velocity in anesthetized children, 830
- see also *Abdallah C*, jt. auth.
Katagai H. See *Kudoh A*, jt. auth.
Kataoka MT. See *Joo HS*, jt. auth.
Kato R, Foëx P. Mycardial protection by anesthetic agents against ischemia-reperfusion injury: an update for anesthesiologists, 777
Katz J. See *Salomons TV*, jt. auth.
Katz SA. See *Moskowitz DM*, jt. auth.
Kavanagh B. See *Skrobik Y*, jt. auth.
Kearney RA, Puchalski SA, Yang HYH, Skakun EN. The inter-rater and intra-rater reliability of a new Canadian oral examination format in anesthesia is fair to good, 232
Keller C. See *Brimacombe J*, jt. auth.
Kelsaka E. See *Sener EB*, jt. auth.
Kennedy B. See *Armstrong KP*, jt. auth.
Kerdilès L. See *Langlois G*, jt. auth.
Kim J. See *Beh T*, jt. auth.
Kitajima T. See *Egawa H*, jt. auth.; *Hamaguchi S*, jt. auth.; *Shinozaki M*, jt. auth.
Klein JJ. See *Moskowitz DM*, jt. auth.
Klein SM. See *Buckenmaier CC III*, jt. auth.
Kligman M, Bruskin A, Sckliamser J, Vered R, Roffman M. Intra-synovial, compared to intra-articular morphine provides better pain relief following knee arthroscopy minisectomy, 380
Kocamanoglu S. See *Sener EB*, jt. auth.
Kohda Y. See *Miyabe M*, jt. auth.
Kohro S, Yamakage M, Takahashi T, Kondo M, Ota K, Namiki A. Intermittent pneumatic compression prevents venous stasis in the lower extremities in the lithotomy position, 144
Kolbitsch C. See *Lorenz IH*, jt. auth.
Kolman J, Morris I. Cricoarytenoid arthritis: a cause of acute upper airway obstruction in rheumatoid arthritis, 729
Kondo M. See *Kohro S*, jt. auth.
Kong S-S. See *Lu C-H*, jt. auth.

Kongsayreepong S, Gunnaleka P, Suraseranivongse S, Pirayavaraporn S, Chowvanayotin S, Montapaneewat T, Manon C. A reusable, custom-made warming blanket prevents core hypothermia during major neonatal surgery, 605
Kudoh A, Katagai H, Takazawa T. Antidepressant treatment for chronic depressed patients should not be discontinued prior to anesthesia, 132
Kusume Y. See *Yokoyama M*, jt. auth.

L
La Coussaye J-E de. See *Lefrant J-Y*, jt. auth.
Ladak S. See *Smith KJ*, jt. auth.
Laha A. See *Manjushree R*, jt. auth.
Lahiri A. See *Manjushree R*, jt. auth.
Lam PKK. See *Hui JKL*, jt. auth.
Lambert AS, Mazer CD, Duke PC. Survey of the members of the cardiovascular section of the Canadian Anesthesiologists' Society on the use of perioperative transesophageal echocardiography - a brief report, 294
Langlois G, Estèbe J-P, Gentili ME, Kerdilès L, Mouilleron P, Ecoffey C. The addition of tramadol to lidocaine does not reduce tourniquet and post-operative pain during *iv* regional anesthesia, 165
Larbig E. See *Piper SN*, jt. auth.
Lass-Flörl C. See *Lorenz IH*, jt. auth.
Laufer B. See *Butler R*, jt. auth.
Laupland KB, Bands CJ. Utility of esophageal Doppler as a minimally invasive hemodynamic monitor: a review, 393
Lazar L. See *Shenkman Z*, jt. auth.
Le Corre P. See *Estèbe J-P*, jt. auth.
Le Verge R. See *Estèbe J-P*, jt. auth.
Lebowitz NE. See *Moskowitz DM*, jt. auth.
Lee A, Ngan Kee WD, Gin T. Prophylactic ephedrine prevents hypotension during spinal anesthesia for Cesarean delivery but does not improve neonatal outcome: a quantitative systematic review, 588
Lee C. See *Ezri T*, jt. auth.
Lee E-J, Sim J-Y, Park J-Y, Hwang J-H, Park P-H, Han S-M. Intrathecal carbachol and clonidine produce a synergistic antiallodynic effect in rats with a nerve ligation injury, 178
Lefrant J-Y, Juan J-M, Bruelle P, Demaria R, Cohendy R, Aya G, Oliva-Lauraire M-C, Peray P, Robert E, de La Coussaye J-E, Eledjam J-J, Dauzat M. Regional blood flows are affected differently by PEEP when the abdomen is open or closed: an experimental rabbit model, 302
Lessard MR. See *Nicole PC*, jt. auth.
Leyland NA. See *Imasogie N*, jt. auth.
Li AH. See *Chung PCH*, jt. auth.
Lienhart A. See *Beaussier M*, jt. auth.
Liepert D. See *Maltby JR*, jt. auth.
Liguoro D. See *Sesay M*, jt. auth.
Lim EHL, Sia ATH, Wong K, Tan HM. Addition of bupivacaine 1.25 mg to fentanyl confers no advantage over fentanyl alone for intrathecal analgesia in early labour, 57
Limoges P. See *Couture P*, jt. auth.
Lin C-H. See *Pang W-W*, jt. auth.
Lin C-J. See *Huang Y-C*, jt. auth.
Lin CC. See *Chung PCH*, jt. auth.
Lin P-L. See *Huang Y-C*, jt. auth.
Ling E, McDonald L, Dinesen TRJ, DuVall D. The OxyArm™ - a new minimal contact oxygen delivery system for mouth or nose breathing, 297
Lingnau W. See *Lorenz IH*, jt. auth.
List WF. See *Dahaba AA*, jt. auth.
Litmanowitz I. See *Shenkman Z*, jt. auth.
Liu K. See *Chia Y-Y*, jt. auth.
Llewellyn R. See *Evans NR*, jt. auth.
Loder RT. See *Munro HM*, jt. auth.
Loeckinger A. See *Brimacombe J*, jt. auth.
Lombroso R. See *Orbach-Zinger S*, jt. auth.
López-Farré A. See *Arriero MM*, jt. auth.
Lorenz IH, Kolbitsch C, Lass-Flörl C, Gritznig I, Vollert B, Lingnau W, Moser PL, Benzer A. Routine handling of propofol prevents contamination as effectively as does strict adherence to the manufacturer's recommendations, 347
Lu C-H, Ho S-T, Kong S-S, Cherng C-H, Wong C-S. Intracranial subdural hematoma after unintended durotomy during spine surgery, 100
Lu M-P. See *Xiao W-J*, jt. auth.
Lucas S. See *Choi PT-L*, jt. auth.
Lucas T. See *Parker RK*, jt. auth.
Luginbuehl I. See *Karsli C*, jt. auth.
Lurie F. See *Jahr JS*, jt. auth.

M
Macarthur A, Halpern S. Best evidence in anesthetic practice (prognosis): the Apgar score predicts 28-day neonatal mortality; commentaries, 46
Mahajan R, Jain K, Batra YK. Submucous fibrosis secondary to chewing of quids: another cause of unanticipated difficult intubation, 309
Mahmoud K. See *Elhakim M*, jt. auth.
Mak PHK, Tsui SL, Ng KFJ. Long-term therapy of chronic non-malignant pain with potent opioids in an active police officer, 575
Makky S. See *Pollard RC*, jt. auth.
Maleck WH. See *Piper SN*, jt. auth.
Maltby JR, Beriault MT, Watson NC, Liepert D, Fick GH. The LMA-ProSeal™ is an effective alternative

to tracheal intubation for laparoscopic cholecystectomy, 857
Malviya S. See *Munro HM*, jt. auth.
Manjushree R, Lahiri A, Ghosh BR, Laha A, Handa K. Intranasal fentanyl provides adequate postoperative analgesia in pediatric patients, 190
Manon C. See *Kongsayreepong S*, jt. auth.
Margolis EJ. See *Moskowitz DM*, jt. auth.
Masawa N. See *Hamaguchi S*, jt. auth.
Massou N. See *Slavov V*, jt. auth.
Maurette P. See *Sesay M*, jt. auth.
Mazer CD. See *Joo HS*, jt. auth.; *Lambert AS*, jt. auth.
Mazer D. See *Skrobik Y*, jt. auth.
McAlister FA. See *Bradley J*, jt. auth.
McDonald L. See *Ling E*, jt. auth.
McFadzean J. See *Pollard RC*, jt. auth.
McKenty S. See *Denault AY*, jt. auth.
McRae K. See *Bradley J*, jt. auth.
McTaggart Cowan RA. See *Archer DP*, jt. auth.
Megnigbeto A. See *Beaussier M*, jt. auth.
Mehran RJ. See *Wherrett CG*, jt. auth.
Mercier FJ. See *Benhamou D*, jt. auth.
Merkel S. See *Munro HM*, jt. auth.
Metais C. See *Park KW*, jt. auth.
Millás I. See *Arriero MM*, jt. auth.
Miller JD. See *Baxter FJ*, jt. auth.
Milne B. See *Aye T*, jt. auth.; *Burjorjee JE*, jt. auth.
Min K. See *Ekatodramis G*, jt. auth.
Minami J. See *Egawa H*, jt. auth.
Miyabe M, Kakiuchi Y, Inomata S, Ohsaka Y, Kohda Y, Toyooka H. Epinephrine does not reduce the plasma concentration of lidocaine during continuous epidural infusion in children, 706
- see also *Nakayama H*, jt. auth.
Mizobuchi S. See *Yokoyama M*, jt. auth.
Montapaneewat T. See *Kongsayreepong S*, jt. auth.
Montecchia F. See *Agrò F*, jt. auth.
Montgomery CJ. See *Pollard RC*, jt. auth.
Montravers P. See *Badaoui R*, jt. auth.
Morgan PJ, Cleave-Hogg D. A worldwide survey of the use of simulation in anesthesia, 659
Morita K. See *Yokoyama M*, jt. auth.
Morley-Forster PK. See *Armstrong KP*, jt. auth.
Morris I. See *Kolman J*, jt. auth.
Moser PL. See *Lorenz IH*, jt. auth.
Moskowitz DM, Perelman SI, Cousineau KM, Klein JJ, Shander A, Margolis EJ, Katz SA, Bennett HL, Lebowitz NE, Ergin MA. Multidisciplinary management of a Jehovah's Witness patient for the removal of a renal cell carcinoma extending into the right atrium, 402
Motamed C, Donati F. Sevoflurane and isoflurane, but not propofol, decrease mivacurium requirements over time, 907
- see also *Slavov V*, jt. auth.
Mouilleron P. See *Langlois G*, jt. auth.
Moulinoux J-P. See *Estève J-P*, jt. auth.
Mowafi HA. See *Al-Metwalli R*, jt. auth.
Mühling J. See *Sablotzki A*, jt. auth.
Muñoz Alameda L. See *Arriero MM*, jt. auth.
Munro HM, Walton SR, Malviya S, Merkel S, Voepel-Lewis T, Loder RT, Farley FA. Low-dose ketorolac improves analgesia and reduces morphine requirements following posterior spinal fusion in adolescents, 461
Murkin J, Hall R. Best evidence in anesthetic practice (prognosis): Cognitive function at hospital discharge predicts long-term cognitive function after coronary artery bypass surgery; commentaries, 497

N

Nafie M. See *Elhakim M*, jt. auth.
Nagatani A, Niiya S, Sumikawa K. High frequency jet ventilation is associated with increased levels of plasma α-atrial natriuretic peptide - a preliminary study, 517
Nakamura M. See *Takenaka I*, jt. auth.
Nakayama H, Takahashi H, Okubo N, Miyabe M, Toyooka H. Xenon and nitrous oxide do not depress cardiac function in an isolated rat heart model, 375
Nakayama M, Ichinose H, Yamamoto S, Kanaya N, Namiki A. The bispectral index response to tracheal intubation is similar in normotensive and hypertensive patients, 458
Namiki A. See *Kohro S*, jt. auth.; *Nakayama M*, jt. auth.; *Yamakage M*, jt. auth.
Narouze S. See *Farag E*, jt. auth.
Neff TA, Fellmann C, Fuechslin RM, Gerbere AC, Weiss M. The Panomat® P-10 micro-volumetric infusion pump is suitable for continuous drug administration at minimal flow rates, 1048
Ng KFJ. See *Mak PHK*, jt. auth.
Ngan Kee WD. See *Lee A*, jt. auth.
Nicole PC, Trépanier CA, Lessard MR. Nausea and vomiting after laparoscopic surgery are not associated with an increased peripheral release of serotonin, 453
Nielsen KC. See *Buckenmaier CC III*, jt. auth.
Niiya S. See *Nagatani A*, jt. auth.

O

O'Brien B, Harmon D, Duggan M, Flynn N. Laryngeal mask cuff inflation at removal does not affect early postoperative laryngopharyngeal morbidity, 871

Oe K. See *Yokoyama M*, jt. auth.
Oettl K. See *Dahaba AA*, jt. auth.
Ofenbartl L. See *Alkaissi A*, jt. auth.
Oh Y-S. See *Park H-P*, jt. auth.
Ohe Y. See *Arakawa M*, jt. auth.
Ohsaka Y. See *Miyabe M*, jt. auth.
Okubo N. See *Nakayama H*, jt. auth.
Okuda Y. See *Egawa H*, jt. auth.; *Hamaguchi S*, jt. auth.; *Shinozaki M*, jt. auth.; *Takamatsu I*, jt. auth.
Oliva-Lauraire M-C. See *Lefrant J-Y*, jt. auth.
Olufolabi AJ. See *Bell EA*, jt. auth.
O'Mara D. See *Duggan M*, jt. auth.
Orbach-Zinger S, Lombroso R, Eidelman LA. Uneventful spinal anesthesia for a patient with carcinoid syndrome managed with long-acting octreotide, 678
Ota K. See *Kohro S*, jt. auth.
Ouendo M. See *Badaoui R*, jt. auth.

P

Pandey CK, Bose N, Dash NR, Singh N, Saxena R. Perioperative management of a patient presenting with a spontaneously ruptured esophagus, 409
Pandey CM. See *Agarwal A*, jt. auth.
Pang W-W, Wu H-S, Lin C-H, Chang D-P, Huang M-H. Metoclopramide decreases emesis but increases sedation in tramadol patient-controlled analgesia, 1029
Park H-P, Baik J-H, Oh Y-S, Ham B-M. Case report: pulmonary soiling after one-lung ventilation with a bronchial blocker, 874
Park J-Y. See *Lee E-J*, jt. auth.
Park KW, Dai HB, Metais C, Comunale ME, Sellke FW. Isoflurane does not further impair microvascular vasomotion in a rat model of subarachnoid hemorrhage, 427
Park P-H. See *Lee E-J*, jt. auth.
Parker RK, Connelly NR, Lucas T, Faheem U, Rizvi AS, El-Mansouri M, Thakkar N, Kamasumadram R, Dixon K, Dunn SM, Gibson C. The addition of hydromorphone to epidural fentanyl does not affect analgesia in early labour, 600
Parlow JL. See *Bracco D*, jt. auth.
Parry N. See *Benhamou D*, jt. auth.
Paut O. See *Bissonnette B*, jt. auth.
Peachey G. See *Denault A*, jt. auth.
Péhourcq F. See *Sesay M*, jt. auth.
Peng P. See *Choyce A*, jt. auth.
Penning DH. See *Bell EA*, jt. auth.
Pennisi MA. See *Cavaliere F*, jt. auth.
Peray P. See *Lefrant J-Y*, jt. auth.
Perelman SI. See *Moskowitz DM*, jt. auth.
Perron R. See *Denault AY*, jt. auth.
Peterson DJ. See *Hudson RJ*, jt. auth.
Petitti T. See *Agrò F*, jt. auth.
Phillips-Bute B. See *Bell EA*, jt. auth.
Pierre S, Benais H, Pouymayou J. Apfel's simplified score may favourably predict patients at risk for postoperative nausea and vomiting, 237
Pinta JC de la. See *Arriero MM*, jt. auth.
Piper SN, Suttner SW, Röhm KD, Maleck WH, Larbig E, Boldt J. Dolasetron, but not metoclopramide prevents nausea and vomiting in patients undergoing laparoscopic cholecystectomy, 1021
Pirayavaraporn S. See *Kongsayreepong S*, jt. auth.
Plante F. See *Denault AY*, jt. auth.
Plaud B. See *Dragne A*, jt. auth.
Pollard RC, Makky S, McFadzean J, Ainsworth L, Goobiec SM, Montgomery CJ. An admixture of 3 $mg \cdot kg^{-1}$ of propofol and 3 $mg \cdot kg^{-1}$ of thiopentone reduces pain on injection in pediatric anesthesia, 1064
Popic PM. See *Yun ES*, jt. auth.
Pouymayou J. See *Pierre S*, jt. auth.
Powles ACP. See *Baxter FJ*, jt. auth.
Pozlep G. See *Vintar N*, jt. auth.
Pradervand Mooser M, Gardaz J-P, Capt H, Spahn DR. Le coût relatif de l'anesthésie pour la cholécystectomie laparoscopique est peu élevé, 540
Prieto-Alvarez P, Tello-Galindo I, Cuenca-Peña J, Rull-Bartomeu M, Gomar-Sancho C. Continuous epidural infusion of racemic methadone results in effective postoperative analgesia and low plasma concentrations, 25
Proietti R. See *Cavaliere F*, jt. auth.; *Conti G*, jt. auth.
Puchalski SA. See *Kearney RA*, jt. auth.

Q

Quaranta S. See *Favaro R*, jt. auth.

R

Rakovec S. See *Vintar N*, jt. auth.
Ramanathan S. See *Vallejo MC*, jt. auth.
Randall J. See *Baxter FJ*, jt. auth.
Rawal N. See *Vintar N*, jt. auth.
Rebufat Y. See *Slavov V*, jt. auth.
Reibnegger G. See *Dahaba AA*, jt. auth.
Reinoso-Barbero F, Saavedra B, Hervilla S, de Vicente J, Tabarés B, Gómez-Criado MS. Lidocaine with fentanyl, compared to morphine, marginally improves postoperative epidural analgesia in children, 67
Rellia P. See *Theodoraki K*, jt. auth.
Reynolds JD. See *Bell EA*, jt. auth.
Rizvi AS. See *Parker RK*, jt. auth.
Robert E. See *Lefrant J-Y*, jt. auth.

Rocchi A, Chung F, Forte L. Canadian survey of post-surgical pain and pain medication experiences, 1053
Roffman M. See *Kligman M*, jt. auth.
Röhm KD. See *Piper SN*, jt. auth.
Rosaeg OP, Campbell N, Crossan ML. Epidural analgesia does not prolong the third stage of labour, 490 - see also *Campbell N*, jt. auth.
Rossaint R. See *de Rossi L*, jt. auth.
Roth W. See *Brimacombe J*, jt. auth.
Rull-Bartomeu M. See *Prieto-Alvarez P*, jt. auth.
Rundshagen I, Schnabel K, Schulte am Esch J. Midlatency auditory evoked potentials do not allow the prediction of recovery from general anesthesia with isoflurane, 361

S

Saavedra B. See *Reinoso-Barbero F*, jt. auth.
Sablotzki A, Czeslick E, Schubert S, Friedrich I, Mühling J, Dehne MG, Grond S, Hentschel T. Iloprost improves hemodynamics in patients with severe chronic cardiac failure and secondary pulmonary hypertension, 1076
Saito Y. See *Yamada Y*, jt. auth.
Sakura S. See *Yamada Y*, jt. auth.
Salomons TV, Wowk AA, Fanning A, Chan VWS, Katz J. Factors associated with refusal to enter a clinical trial: epidural anesthesia is a deterrent to participation, 583
Sarihasan B. See *Sener EB*, jt. auth.
Saulys A. See *Yun ES*, jt. auth.
Sawchuk C. See *Fayad A*, jt. auth.
Saxena R. See *Pandey CK*, jt. auth.
Schmid ER. See *Bettex DA*, jt. auth.
Schmidlin D. See *Bettex DA*, jt. auth.
Schnabel K. See *Rundshagen I*, jt. auth.
Schubert S. See *Sablotzki A*, jt. auth.
Schulte am Esch J. See *Rundshagen I*, jt. auth.
Schulz G. See *Weiss M*, jt. auth.
Sckliamser J. See *Kligman M*, jt. auth.
Sellke FW. See *Park KW*, jt. auth.
Sener EB, Sarihasan B, Ustun E, Kocamanoglu S, Kelsaka E, Tur A. Awake tracheal intubation through the intubating laryngeal mask airway in a patient with halo traction, 610
Sesay M, Dousset V, Liguoro D, Péhourcq F, Caillé J-M, Maurette P. Intraosseous lidocaine provides effective analgesia for percutaneous vertebroplasty of osteoporotic fractures, 137
Sesay M, Vignes J-R, Liguoro D, Crozat P, Boulard G, Guerin J, Barat M, Maurette P. L'hyperréflexie autonome induite par la stimulation des racines sacrées est détectée par l'analyse spectrale de l'ECG, 936
Sfyrakis P. See *Theodoraki K*, jt. auth.
Shander A. See *Moskowitz DM*, jt. auth.
Shenkman Z, Hoppenstein D, Litmanowitz I, Shorer S, Gutermacher M, Lazar L, Erez I, Jedeikin R, Freud E. Spinal anesthesia in 62 premature, former-premature or young infants - technical aspects and pitfalls, 262
Sheridan P. See *Couture P*, jt. auth.; *Taillefer J*, jt. auth.
Shinozaki M, Usui Y, Yamaguchi S, Okuda Y, Kitajima T. Recovery of psychomotor function after propofol sedation is prolonged in the elderly, 927
Shorer S. See *Shenkman Z*, jt. auth.
Shrestha BM, Singh BB, Gautam MP, Chand MB. The oxygen concentrator is a suitable alternative to oxygen cylinders in Nepal, 8
Sia ATH. See *Lim EHL*, jt. auth.
Siddik-Sayyid S. See *Baraka A*, jt. auth.
Sim J-Y. See *Lee E-J*, jt. auth.
Singh B, Bhardwaj V. Continuous mandibular nerve block for pain relief. A report of two cases, 951
Singh BB. See *Shrestha BM*, jt. auth.
Singh D. See *Agarwal A*, jt. auth.
Singh K. See *Hudson RJ*, jt. auth.
Singh N. See *Pandey CK*, jt. auth.
Singh PK. See *Agarwal A*, jt. auth.
Singh U. See *Agarwal A*, jt. auth.
Skakun EN. See *Kearney RA*, jt. auth.
Skrobik Y, Kavanagh B, Mazer D, Cook D. Toronto critical care medicine symposium report, 513
Skrobik YK, Filep JG. Epi and endomyocardial pH allows the detection of acute right ventricular ischemia in pigs: a new evaluation method, 84
Slavov V, Motamed C, Massou N, Rebufat Y, Duvaldestin P. Systolic blood pressure, not BIS, is associated with movement during laryngoscopy and intubation, 918
Smith CE, Boyer D. Cricoid pressure decreases ease of tracheal intubation using fibreoptic laryngoscopy (WuScope System™), 614
Smith KJ, Ladak S, Choi PT-L, Dobranowski J. The cricoid cartilage and the esophagus are not aligned in close to half of adult patients, 503
Song D. See *Wong J*, jt. auth.
Spahn DR. See *Pradervand Mooser M*, jt. auth.
Spielman FJ. Complete separation of the tube from the mask during removal of a disposable laryngeal mask airway, 990
Splinter W. See *Beh T*, jt. auth.
Stacey MRW. See *Williams DJ*, jt. auth.
Steele SM. See *Buckenmaier CC III*, jt. auth.
Sumikawa K. See *Nagatani A*, jt. auth.
Sun W-Z. See *Huang Y-C*, jt. auth.
Suraseranivongse S. See *Kongsayreepong S*, jt. auth.

Sutherland GR. See *Archer DP*, jt. auth.
Suttner SW. See *Piper SN*, jt. auth.
Swenerton JE. See *Douglas MJ*, jt. auth.
Sykes J. See *Thompson K*, jt. auth.

T
Tabarés B. See *Reinoso-Barbero F*, jt. auth.
Taillefer J, Couture P, Sheridan P, Girard A, Babin D, Denault A. A comprehensive strategy to avoid transesophageal echocardiography probe damage, 500
Takahashi H. See *Nakayama H*, jt. auth.
Takahashi T. See *Kohro S*, jt. auth.
Takamatsu I, Karasawa F, Okuda T. Dopamine may preserve the myocardial oxygen balance better than dobutamine when administered with milrinone, 968
Takazawa T. See *Kudoh A*, jt. auth.
Takenaka I, Aoyama K, Nakamura M, Fukuyama H, Urakami Y, Takenaka Y, Kadoya T. Oral styletted intubation under video control in a patient with a large mobile glottic tumour and a difficult airway, 203
Takenaka Y. See *Takenaka I*, jt. auth.
Takeuchi L. See *Choi PT-L*, jt. auth.
Tan HM. See *Lim EHL*, jt. auth.
Tang G-Z. See *Xiao W-J*, jt. auth.
Técsy M. See *Benhamou D*, jt. auth.
Tello-Galindo I. See *Prieto-Alvarez P*, jt. auth.
Thakkar N. See *Parker RK*, jt. auth.
Thanopoulos A. See *Theodoraki K*, jt. auth.
Theodoraki K, Rellia P, Thanopoulos A, Tsourelis L, Zarkalis D, Sfyrakis P, Antoniou T. Inhaled iloprost controls pulmonary hypertension after cardiopulmonary bypass, 963
Thompson K, Wisenberg G, Sykes J, Thompson RT. Similar long-term cardiovascular effects of propofol or isoflurane anesthesia during ischemia/reperfusion in dogs, 978
Thompson RT. See *Thompson K*, jt. auth.
Thomson IR. See *Hudson RJ*, jt. auth.
Tilleul P. See *Beaussier M*, jt. auth.
To T. See *Jacka MJ*, jt. auth.
Tome J. See *Farag E*, jt. auth.
Tordiglione P. See *Favaro R*, jt. auth.
Tormey W. See *Duggan M*, jt. auth.
Toyooka H. See *Fujii Y*, jt. auth.; *Miyabe M*, jt. auth.; *Nakayama H*, jt. auth.
Trépanier CA. See *Nicole PC*, jt. auth.
Truman C. See *Johnson D*, jt. auth.
Tsai S-K. See *Huang Y-C*, jt. auth.
Tsai S-Y. See *Chang H*, jt. auth.
Tsourelis L. See *Theodoraki K*, jt. auth.
Tsui SL. See *Mak PHK*, jt. auth.
Tsujiguchi N. See *Yamakage M*, jt. auth.
Tur A. See *Sener EB*, jt. auth.

U
Urakami Y. See *Takenaka I*, jt. auth.
Ustun E. See *Sener EB*, jt. auth.
Usui Y. See *Shinozaki M*, jt. auth.

V
Vallejo MC, Ramanathan S. Plasma lidocaine concentrations are higher in twin compared to singleton newborns following epidural anesthesia for Cesarean delivery, 701
VanDenKerkhof EG, Goldstein DH, Wilson R. A survey of directors of Canadian academic acute pain management services: the nursing team members role - a brief report, 579
Varin F. See *Dragne A*, jt. auth.
Vered R. See *Kligman M*, jt. auth.
Vignes J-R. See *Sesay M*, jt. auth.
Vintar N, Pozlep G, Rawal N, Godec M, Rakovec S. Incisional self-administration of bupivacaine or ropivacaine provides effective analgesia after inguinal hernia repair, 481
Voepel-Lewis T. See *Munro HM*, jt. auth.
Vohra A. See *Ahmed MZ*, jt. auth.
Vollert B. See *Lorenz IH*, jt. auth.
von Klobucar F. See *Dahaba AA*, jt. auth.

W
Walton SR. See *Munro HM*, jt. auth.
Wang M-H. See *Huang Y-C*, jt. auth.
Wang P-N. See *Chia Y-Y*, jt. auth.
Watson JT. See *Armstrong KP*, jt. auth.
Watson NC. See *Maltby JR*, jt. auth.
Weiss M, Schulz G, Fasnacht M, Balmer C, Fischer JE, Gerber AC, Bucher H-U, Baenziger O. Transcutaneously measured near-infrared spectroscopic liver tissue oxygenation does not correlate with hepatic venous oxygenation in children, 824 - see also *Dullenkopf A*, jt. auth.; *Neff TA*, jt. auth.
Wennberg R. See *Jenkins K*, jt. auth.
Wherrett CG, Mehran RJ, Beaulieu M-A. Cerebral arterial gas embolism following diagnostic bronchoscopy: delayed treatment with hyperbaric oxygen, 96
White H, Baker A. Continuous jugular venous oximetry in the neurointensive care unit - a brief review, 623
Williams DJ, Stacey MRW. Rapid and complete occlusion of a heat and moisture exchange filter by pulmonary edema (clinical report), 126
Wilson R. See *VanDenKerkhof EG*, jt. auth.
Wisenberg G. See *Thompson K*, jt. auth.
Wong C-S, Chang Y-C, Yeh C-C, Huang G-S, Cherng C-H. Loss of intrathecal morphine analgesia in terminal cancer patients is associated with high levels of excitatory amino acids in the CSF, 561

- see also *Lu C-H*, jt. auth.
Wong DT. See *Jenkins K*, jt. auth.
Wong J, Song D, Blanshard H, Grady D, Chung F. Titration of isoflurane using BIS index improves early recovery of elderly patients undergoing orthopedic surgeries, 13
Wong K. See *Lim EHL*, jt. auth.
Wowk AA. See *Salomons TV*, jt. auth.
Wu H-S. See *Pang W-W*, jt. auth.

X

Xiao W-J, Deng X-M, Tang G-Z, Lu M-P, Xu K-L. Caudal anesthesia reduces the minimum alveolar concentration of enflurane for laryngeal mask airway removal in boys, 194
Xu K-L. See *Xiao W-J*, jt. auth.

Y

Yaacoub C. See *Baraka A*, jt. auth.
Yamada Y, Doi K, Sakura S, Saito Y. Anesthetic management for a patient with Jansky-Bielschowsky disease, 81
Yamaguchi S. See *Shinozaki M*, jt. auth.
Yamakage M, Namiki A. Calcium channels - basic aspects of their structure, function and gene encoding; anesthetic action on the channels - a review, 151
Yamakage M, Tsujiguchi N, Chen X, Kamada Y, Namiki A. Sevoflurane inhibits contraction of uterine smooth muscle from pregnant rats similarly to halothane and isoflurane, 62
- see also *Kohro S*, jt. auth.
Yamamoto S. See *Nakayama M*, jt. auth.
Yang H. See *Fayad A*, jt. auth.
Yang HYH. See *Kearney RA*, jt. auth.
Yang MW. See *Chung PCH*, jt. auth.
Yee I. See *Armstrong KP*, jt. auth.
Yeh C-C. See *Wong C-S*, jt. auth.
Yokoyama M, Itano Y, Kusume Y, Oe K, Mizobuchi S, Morita K. Total spinal anesthesia provides transient relief of intractable pain, 810
Yun ES, Saulys A, Popic PM, Arndt GA. Single-lung ventilation in a pediatric patient using a pediatric fibreoptically-directed wire-guided endobronchial blocker, 256

Z

Zarkalis D. See *Theodoraki K*, jt. auth.
Zito G. See *Conti G*, jt. auth.